

SIRA and scattering in the IPM Appendix

T. S. Bastian
NRAO



Dayton Jones kindly provided me with a sample ALFA configuration, a distribution of 16 antennas in a spherical shell of 100 km diameter.

- o As a zeroeth order exercise, I produced 5 scaled versions of Dayton's array. The array diameters were 2, 5, 10, 20, and 50 km.

- o Snapshot uv data bases were then produced for each array at a frequency of 1 MHz.

- o A source model was passed through the instrument. The model was a simple Gaussian with a FWHM angular size of 5 degrees (consistent with the ISEE-3 results of Steinberg et al 1985 and my simple PWE calculations).

- o The source flux was arbitrarily chosen to be 1 Jy.

- o Note: the data were noiseless.

The result, briefly, is as follows:

As can be seen, the 5 km array is best matched to the source size. The 2 km under-resolves the source while the 10 km array over-resolves the source. The 20 and 50 km arrays grossly over-resolve the source.

Comments:

It may be possible to use elongated configurations that can satisfy both solar and astrophysical mapping requirements, but given the fact that solar source mapping must be supported to large solar elongations (90°?) this must be studied carefully.

An alternative is that the array can be reconfigured in time, beginning with a compact configuration and slowly moving to a larger configuration.

Regardless of the solution adopted, these simple calculations are meant to be illustrative only. Much, much more work needs to be done along these lines.


- o Model (projected) uv coverage for a spherical array 50 km in diameter.

- o In the uv domain, it samples baselines up to 166λ .

- o Approximately uniform coverage is apparent.


- o All other arrays are scaled versions of this one.

- o The most compact is 2 km, yielding a maximum baseline of only ~ 6 wavelengths.




2 km array

Visibility amplitude
as a function of uv
distance. Recall
that the total
source flux is 1 Jy.




2 km array

Dirty map (i.e., no deconvolution) of the model source.




5 km array

Visibility amplitude
as a function of uv
distance.




5 km array

Dirty map of the
model source.



10 km array

Visibility amplitude
as a function of uv
distance.




10 km array

Dirty map of the
model source.

Note that the
sidelobe level is as
high as 50%.

A significant
amount of source
flux has been
resolved out.




20 km array

Visibility amplitude
as a function of uv
distance.

Only ~10% of the
baselines see
correlated flux.

The source is
grossly over-
resolved.




20 km array

Dirty map of the
model source.

Without prior
knowledge, the
source is largely
indistinguishable
from the sidelobes.


Most of the source
flux is resolved
out.



50 km array

Visibility amplitude
as a function of uv
distance.

There is no
correlated flux
seen on any
baseline, only
(numerical) noise.



50 km array

Dirty map of the
model source.

No source is
present, just
numerical noise.

